

Bed Wetting—Silent Suffering: An Approach to Enuresis and Voiding Disorders in Children: Correspondence

Narendra Rathi · Manisha Rathi

Received: 6 November 2013 / Accepted: 12 December 2013
© Dr. K C Chaudhuri Foundation 2014

To the Editor: I congratulate authors for informative and practical article on Enuresis [1]. I seek clarification regarding a couple of issues.

1. It is advised to compare in Frequency Volume Diary the volume of urine voided each time to expected normal bladder volume for respective age of the child. What are the lower limits of urinary volume for various ages to define small bladder capacity?
2. It is mentioned that smaller than expected bladder capacity are likely to be Desmopressin resistant, but Desmopressin with alarm is advised for children with excessive urine output and reduced bladder capacity.

Contribution Dr. Narendra Rathi will act as guarantor for the paper.

Conflict of Interest None.

Role of Funding Source None.

Reference

1. Kanitkar M, Ramamurthy HR. Bed wetting-silent suffering: An approach to enuresis and voiding disorders in children. Indian J Pediatr. 2013;80:750–3.

N. Rathi (✉) · M. Rathi
Department of Pediatrics, Rathi Children Hospital, Civil Lines,
Akola 444001, Maharashtra, India
e-mail: drnbrathi@hotmail.com